

TOM J. POUSTI, MD, F.A.C.S.
PLASTIC AND RECONSTRUCTION SURGERY

INFORMED CONSENT FOR BREAST AUGMENTATION SURGERY

INSTRUCTIONS

This is an informed consent document that has been prepared to help inform you about augmentation mammoplasty, its risks, and alternative treatments.

It is important that you read this information carefully and completely. Please initial each page, indicating you have read the page and sign the consent for surgery as proposed by your plastic surgeon.

GENERAL INFORMATION

Augmentation Mammoplasty is a surgical operation performed to enlarge the breasts for a number of reasons:

- To enhance the body contour of a woman, who for personal reasons feels that her breast size is too small.
- To correct a loss in breast volume after pregnancy.
- To balance breast size, when there exists a significant difference between the size of the breasts.

As a reconstructive technique for various conditions.

Replacement of breast implants for medical or cosmetic reasons.

The shape and size of the breasts prior to surgery will influence both the recommended treatment and the final results. If the breasts are not the same size or shape before surgery, it is unlikely that they will be completely symmetrical afterward.

Silicone gel filled implants have been approved by the U.S. Food and Drug Administration (FDA) as of November 17th,2006. Breast enlargement is accomplished by inserting a breast implant either behind the breast tissue or under the chest muscles. Incisions are made to keep scars as inconspicuous as possible, usually under the breast, around the lower pole of the areola, or in the armpit. The method of inserting and positioning breast implants will depend on your preferences, your anatomy and your surgeon's recommendation.

Patients undergoing augmentation mammoplasty surgery must consider the possibility of future revisionary surgery. Breast implants cannot be expected to last forever.

ALTERNATIVE TREATMENT

Augmentation mammoplasty is an elective surgical operation. Alternative treatment would consist of the use of external breast prostheses or padding, or the transfer of other body tissues to enlarge breast size.

RISKS OF AUGMENTATION MAMMOPLASTY SURGERY

Every surgical procedure involves a certain amount of risk and it is important that you understand the risks involved with augmentation mammoplasty. Additional information concerning breast implants may be obtained from the FDA, package inserts sheets supplied by the implant manufacturer, or other information pamphlets required by individual state laws.

An individual's choice to undergo a surgical procedure is based on the comparison of the risk to potential benefit. Although the majority of women do not experience the following complications, you should discuss each of them with your plastic surgeon to make sure you understand the risks, potential complications, and consequences of breast augmentation.

Bleeding

It is possible, though unusual, to experience a bleeding episode during or after surgery. Should post-operative bleeding occur, it might require emergency treatment to drain accumulated blood (hematoma). Do not take aspirin or anti-inflammatory medications for ten days before surgery, as this may increase the risk of bleeding.

Infection

Infection is unusual after this type of surgery. It may appear in the immediate postoperative period or at any time following the insertion of a breast implant. Subacute or chronic infections may be difficult to diagnose. Should an infection occur, treatment including antibiotics, possible removal of the implant, or additional surgery may be necessary. Infections with the presence of a breast implant are harder to treat than infections in normal body tissue. If an infection does not respond to antibiotics, the breast implant may have to be removed. After the infection is treated, a new breast implant can usually be reinserted. It is extremely rare that an infection would occur around an implant from a bacterial infection elsewhere in the body, however, prophylactic antibiotics may be considered for subsequent dental or other surgical procedures.

Capsular Contracture

Scar tissue, which forms internally around the breast implant, can tighten around the breast implant, can tighten and make the breast firm and possibly painful. Excessive firmness of the breasts can occur soon after surgery or years later. Though the occurrence of symptomatic capsular contracture is not predictable, it generally occurs in less than 20 percent of patients. The incidence of symptomatic capsular contracture can be expected to increase over time. Capsular contracture may occur on one side, both sides, or not at all. Treatment for capsular contracture may require surgical implant replacement, or implant removal.

Change in nipple and skin sensation

Some change in nipple sensation is not unusual right after surgery. After several months, most patients have normal sensation. Partial or permanent loss of nipple and skin sensation may occur occasionally.

Skin Scarring

Excessive scarring is uncommon. In rare cases, abnormal scars may result. Scars may be unattractive and of different color than surrounding skin. Additional surgery may be needed to treat abnormal scarring after surgery.

Implant Failure

Breast implants, similar to other medical devices, can fail. Implants can break or leak. When a saline filled implant deflates, its salt-water filling will be absorbed by the body. Rupture can result of an injury, inapparent cause, or during mammography. It is possible to damage an implant at the time of surgery. Damaged or broken implants cannot be repaired. Ruptured or deflated implants require replacement or removal. Breast implants cannot be expected to last forever.

Degradation of breast implants

It is possible that small pieces of the implant material may separate from the outer surface of breast implants. This is of unknown significance.

Implant Extrusion

Lack of adequate tissue coverage or infection may result in exposure and extrusion of the implant. Skin breakdown has been reported with the use of steroid drugs or after radiation therapy to breast tissue. If tissue breakdown occurs and the implant becomes exposed, implant removal may be necessary. Smoking may interfere with the healing process.

Mammography

Breast implants may make mammography more difficult and may obscure the detection of breast cancer. Implant rupture can occur from breast compression during mammography. Inform your mammography technologist of the presence of breast implants so that appropriated mammogram studies may be obtained. Patients with capsular contracture may find mammogram techniques painful and the difficulty of breast imaging will increase with the extent of contracture. Ultrasound, specialized mammography and MRI studies may be of benefit to evaluate breast lumps and the condition of the implants. Because more x-ray views are necessary with specialized mammography techniques, women with breast implants will receive more radiation than women without implants who receive a normal exam. However, the benefit of the mammogram in finding cancer outweighs the risk of additional x-rays.

Skin Wrinkling and Rippling

Visible and palpable wrinkling of implants can occur. Some wrinkling is normal and expected. This may be more pronounced in patients who have saline filled implants or thin breast tissue. It may be possible to feel the implant valve. Some patients may find palpable valve and wrinkles cosmetically undesirable. Palpable wrinkling and/or folds may be confused with palpable tumors and questionable cases must be investigated. An implant may become visible at the surface of the breast as a result of the device pushing through layers of skin.

Pregnancy and breast feeding

Although many women with breast implants have successfully breast fed their babies, it is not known if there are increased risks in nursing for a woman with breast implants or if the children of women with breast implants are more likely to have health problems. There is insufficient evidence regarding the absolute safety of breast implants in relation to fertility, pregnancy, or breast feeding. Some women with breast implants have reported health problems in their breast fed children. Only very limited research has been conducted in this area and at this time there is no scientific evidence that there is a problem.

Asymmetry

Many women have naturally asymmetrical breasts. Although surgery will attempt to correct the difference, asymmetry may continue to be present following surgery. Should this occur, additional surgery might be necessary to improve the differences. Additional costs may occur.

Calcification

Calcium deposits can form in the scar tissue surrounding the implant and may cause pain, firmness, and be visible on mammography. These deposits must be identified as different from calcium deposits that are a sign of breast cancer. Should this occur, additional surgery might be necessary to remove and examining calcifications.

Implant Displacement

Displacement or migration of breast implant may occur from its initial placement and can be accompanied by discomfort and/or distortion in breast shape. Difficult techniques of implant placement may increase the risk of displacement or migration. Additional surgery may be needed to correct this problem.

Surface Contamination of Implants

Skin oil, lint from surgical drapes, or talc may become deposited on the surface of the implant at the time of insertion. The consequence of this is unknown.

Surgical Anesthesia

Both local and general anesthesia involve risk. There is the possibility of complications, injury, and even death from all forms of surgical anesthesia or sedation.

Chest wall Deformity

Chest wall deformity has been reported secondary to the use of tissue expanders and breast implants. The consequences of chest wall deformity is of unknown significance.

Unusual activities and occupations

Activities and occupations which have the potential for trauma to the breast could potentially break or damage breast implants, or cause bleeding.

Allergic reactions

In rare cases, local allergies to tape, suture material, or topical preparations have been reported. Systemic reactions, which are more serious, may result from drugs used during surgery and prescription medicines. Allergic reactions may require additional treatment.

Breast Disease

Current medical information does not demonstrate an increased risk of breast disease or breast cancer in women who have breast implant surgery for either cosmetic or reconstructive purposes. Breast disease can occur independently of breast implants. It is recommended that all women perform self examination of their breasts, have mammography according to the American Cancer Society guidelines, and seek professional care should they notice a breast lump.

Seroma

Fluid may accumulate around the implant following surgery, trauma or vigorous exercise. Additional treatment may be necessary to drain fluid accumulation around breast implants.

Long Term Results

Subsequent alterations in breast shape may occur as the result of aging, weight loss or gain, pregnancy, or other circumstances not related to augmentation mammoplasty. Breast sagging may occur normally.

Thrombosed Veins

Thrombosed veins, which resemble cords occasionally develop in the area of the breast and resolve without medical or surgical treatment.

Immune System Diseases and Unknown Risks

Some women with breast implants have reported symptoms similar to those of known diseases of the immune system such as systemic lupus erythematosus, rheumatoid arthritis, scleroderma, and other arthritis like conditions. A connection between implanted silicone and connective tissue disorders has been reported in the medical literature. To date, there is no scientific evidence that women with either silicone gel filled or saline filled breast implants have an increased risk of these diseases, but the possibility cannot be excluded. If a causal relationship is established, the theoretical risk of immune and unknown disorders may be low. The effects of breast implants in individuals with pre-existing connective tissue disorders is unknown.

Unlike silicone gel filled implants, the saline filled implants contain salt water. Any risk related to silicone gel would not be associated with saline filled implants. However, gel filled and saline filled devices have a silicone rubber envelope. An increased risk of autoimmune disease is possible even from saline implants. Reliable medical laboratory tests to determine antibodies to silicone do not exist. It has not been proved that there is a relationship between silicone antibodies and disease in women with breast implants. Currently, there is insufficient evidence to state that there is a health benefit from removing either breast implants and scar tissue capsules or that removal will alter autoimmune disease or prevent its potential occurrence.

In very few women who have breast implants, a variety of other symptoms and conditions have been reported, suggestive of an autoimmune multiple-sclerosis-like syndrome. Additional complaints involve the musculoskeletal, skin, nervous, and immune systems. The relationship of breast implants to these conditions has been hypothesized, although not scientifically proven. Because such disease states are rare, they are difficult to research.

Current studies have only looked for the symptoms of known autoimmune diseases, rather than the variety of symptoms that women report experiencing. Some of the reported symptoms include:

- ◆ swelling or joint pain or arthritis like pain
- ◆ general aching
- ◆ unusual hair loss
- ◆ unexplained or unusual loss of energy
- ◆ greater chance of getting colds, viruses, flu
- ◆ swollen glands or lymph nodes
- ◆ rash
- ◆ memory problems, headaches
- ◆ muscle weakness or burning
- ◆ nausea, vomiting
- ◆ irritable bowel syndrome
- ◆ fever

Questions have been raised about the potential for the saline solution used to fill breast implants to become contaminated with bacteria or fungus. These organisms may present a risk to the patient in the event of implant leakage or deflation.

There is the possibility of unknown risks associated with silicone breast implants and tissue expanders.

Toxic shock syndrome – This is an extremely rare complication following breast augmentation, reconstruction, or tissue expansion with silicone implants.

Unsatisfactory result – You may be disappointed with the results of surgery. Asymmetry in implant placement, breast shape and size may occur after surgery. Unsatisfactory surgical scar location or displacement may occur. Pain may occur following surgery. It may be necessary to perform additional surgery to improve your results.

Removal/replacement of breast implants – Future removal or replacement of breast implants and the surrounding scar tissue envelope involves a surgical procedure with risks and potential complications.

HEALTH INSURANCE

Most health insurance companies exclude coverage for cosmetic surgical operations such as the augmentation mammoplasty and any complications that might occur from surgery. Some insurance carriers may possibly exclude breast diseases in patients who have breast implants. Please carefully review your health insurance subscriber information pamphlet.

ADDITIONAL SURGERY NECESSARY

Should complications occur, additional surgery or other treatments might be necessary. Even though risks and complications occur infrequently, the risks cited are particularly associated with augmentation mammoplasty; other complications and risks can occur but are even more uncommon. The practice of medicine and surgery is not an exact science. Although good results are expected, there is no guarantee or warranty expressed or implied on the results that may be obtained.

FINANCIAL RESPONSIBILITIES

The cost of surgery involves several charges for the services provided. The total includes fees charged by your doctor, the cost of implants, and surgical supplies, anesthesia, laboratory tests, and possible outpatient hospital charges, depending on where the surgery is performed. Depending on whether the cost of surgery is covered by an insurance plan you will be responsible for necessary co-payments, deductibles, and charges not covered. Additional costs may occur should complications develop from the surgery. Secondary surgery or hospital day surgery charges involved with revisionary surgery would also be your responsibility.

DISCLAIMER

Informed consent documents are used to communicate information about the proposed surgical treatment of a disease or condition along with disclosure of risks and alternative forms of treatment. The informed consent process attempts to define principles of risk disclosure that should generally meet the needs of most patients in most circumstances.

However, informed consent documents should not be considered all inclusive in defining other methods of care and risks encountered. Your plastic surgeon may provide you with additional or different information which is based on all the facts in your particular case and the state of medical knowledge.

Informed consent documents are not intended to define or serve as the standard of medical care. Standards of medical care are determined on the basis of all of the facts involved in an individual case and are subject to change as scientific knowledge and technology advance and as practice patterns evolve.

I have been informed of the Mentor Enhanced Advantage Limited Warranty. I understand that, should I elect to purchase this program, I must do so within 30 days of my qualifying implantation. I also understand the steps necessary to purchase the program as outlined in the Mentor Advantage Limited Warranty brochure.

It is important that you read the above information carefully and have all of your questions answered before signing the consent on the next page.

CONSENT FOR SURGERY/PROCEDURE or TREATMENT

I hereby authorize Dr. Tom J. Pousti and such assistant as may be selected to perform the following procedure or treatment:

BREAST AUGMENTATION

I have received the following information sheet:

INFORMED CONSENT FOR AUGMENTATION MAMMOPLASTY

I recognize that during the course of the operation and medical treatment or anesthesia, unforeseen conditions may necessitate different procedures than those above. I therefore authorize the above physician and assistants of designees to perform such other procedures that are in the exercise of his or her professional judgment necessary and desirable. The authority granted under this paragraph shall include all conditions that require treatment and are not known to my physician at the time the procedure has begun.

I consent to the administration of such anesthetics considered necessary or advisable. I understand that all forms of anesthesia involve risk and possibility of complications, injury, and sometimes death.

I acknowledge that no guarantee has been given by anyone as to the results that may be obtained.

I consent to the photographing or televising of the operation or procedure to be performed, including appropriate portions of my body, for medical, scientific or educational purposes, provided my identity is not revealed by the pictures.

For purposes of advancing medical education, I consent to the admittance of observers to the operating room.

I consent to the disposal of any tissue, medical devices or body parts, which may be removed.

I authorize the release of my social security number to appropriate agencies for legal reporting and medical device registration if applicable.

IT HAS BEEN EXPLAINED TO ME IN A WAY THAT I UNDERSTAND:

- a) THE ABOVE TREATMENT OR PROCEDURE TO BE UNDERTAKEN
- b) THERE MAY BE ALTERNATIVE PROCEDURE OR METHODS OF TREATMENT
- c) THERE ARE RISKS TO THE PROCEDURE OR TREATMENT PROPOSED

Sign A or B Below

A. I consent to the treatment or procedure and the above listed items. I have been asked if I want more detailed explanation, but I am satisfied with the explanation and do not want more information.

Patient or person authorized to sign for patient

Date

B. I consent to the treatment or procedure and above listed items. I requested and received, in substantial detail, further explanation of the procedure or treatment, other alternative procedures or methods or treatment and information about the material risks or the procedure of treatment.

Patient or person authorized to sign for patient

Date